

MGI Michiana Gastroenterology Inc.

Telephone: (574) 234-0049

Fax: (574) 234-0053

www.michianagastro.com

Colonoscopy Instructions

with GOLYTELY/NULYTELY/TRILYTE PREP

Please be aware, there will be 2 separate charges for this procedure. One for the Facility and one for the Doctor

Getting Ready for your Colonoscopy Please read today!

Patient: _____

Dr. _____

Test Date: ____/____/____

Arrival Time: _____ am/pm

Procedure Time: _____ am/pm

_____ Michiana Endoscopy Center
53830 Generations Drive

_____ St. Joseph's Medical Center
Main Entrance Registration

_____ Memorial Hospital
Patient Registration – Main Entrance

If you need to cancel or reschedule your test please call the office at 574-234-0049.

Do this 7 days before your procedure:

- Stop iron medication and iron supplements.

Do these things 4 days before your test:

- Arrange to have someone drive you home. You will receive medicine to keep you relaxed during your procedure, so you cannot drive home or take a cab home alone.
- Stop taking aspirin. It is okay to take Tylenol (Acetaminophen).
- If you take Coumadin, check with the doctor who ordered it to see if you can stop taking it for the 4 days before the test.
- If you take insulin call the doctor who ordered it for any adjustments you need to make the day of prep and the test day. We do not want your blood sugar to be too low.

Do these things 1 day before your test (Prep Day):

- DO NOT EAT ANY SOLID FOOD TODAY**
- You may only have clear liquids all day and night. **NO RED OR PURPLE.** Clear liquids are: tea or coffee (no milk), broth, apple juice, white cranberry juice, 7 up, ginger ale and water. You may also have Gatorade, popsicles and Jell-O, just remember no red or purple.
- Do not drink alcohol.
- Start drinking your prep according to your doctor's instructions, which are located on the back of this page.
 - Note: Some people say taking sips of broth or sucking on a lime in between glasses of prep help to clear the taste.
 - If you feel too full, drink the prep every 30 minutes.
- Keep drinking clear liquids for the rest of the day. This will help the prep work better for you.
- The prep you are drinking will make you go to the bathroom many times. This is needed so your colon will be cleaned out for the procedure.
- You may use Vaseline, any diaper rash cream or baby wipes, etc...on your bottom. This will help it not get so sore.

Do these things on the day of your test: DO NOT EAT ANY SOLID FOOD

- You may continue drinking clear liquids, but stop drinking 2 hours before your test.
- Take your usual morning pills except for Coumadin and Aspirin.
- DO NOT take your diabetic pill(s).
- If you usually check your blood sugar, check it this morning.
- Bring the person who is driving you home with you to the Endoscopy Center.
- Your arrival time to your discharge time is about 2 ½ hours.
- Bring insurance card, photo id and all medications to your appointment.

OVER →

Day Before Examination

FOLLOW PRINTED INSTRUCTIONS ON THIS SHEET NOT THE COLON PREP PACKAGING.

1. Upon waking, immediately take the four (4) Dulcolax Tablets all at once with 8 ounces of clear liquid.
2. **Drink only clear liquids for breakfast, lunch and dinner.** Clear liquids throughout the day are encouraged.
3. In the morning, add lukewarm water to the top line on the container. Mix it to dissolve the powder. It is sometimes easier to drink the prep through a straw and/or chilled. If you choose you may put the container in the refrigerator to chill.
4. **You will be taking a laxative and it will give you diarrhea. Stay close to the restroom.**
5. **12:00 PM** begin drinking Golytely/Nulytely/Trilyte Prep (you may start earlier).
You will be drinking the entire container of the Golytely/Nulytely/Trilyte Prep today. Drink 1(8 oz) glass of the mixture every 15 minutes.
6. It is important to drink at least eight (8) glasses of clear liquids throughout the day to prevent dehydration. Choose from the above list.
7. If you have rectal discomfort from the prep, Tucs, A and D ointment or Desitin ointment is recommended for use around the rectal area.
8. If you awaken during the night with brown or solid stool drink more clear liquids. Tell the nurse at the endoscopy center if you are still having brown or solid stool upon arrival.

Day of Examination

1. **DO NOT EAT ANY SOLID FOOD until after your procedure.**
2. DRINKING LIQUIDS ARE ENCOURAGED TO PREVENT DEHYDRATION. YOU MAY DRINK LIQUIDS UP TO 2 HOURS BEFORE YOUR EXAMINATION.
3. **REMINDER:** You must be accompanied by a driver because you will be sedated for the procedure. Do not plan to return to work or make any important decisions the rest of the day.

Phone Numbers if there are questions:

Michiana Gastroenterology, Inc.
Hours of operation

(574) 234-0049

Monday, Tuesday and Thursday 8:00am – 5:00pm

Wednesday & Friday 8:00am – 1:30pm

Michiana Endoscopy Center
Hours of operation

(574) 271-0893 or 1-866-669-8743 (toll free)

Monday – Thursday 5:45am – 5:00 pm

Friday 5:45am – 4:00pm

St. Joseph Regional Medical Center
Memorial Hospital

(574) 237-7736 8am – 4pm

(574) 647-6700 8am – 4pm

IN CASE OF AN EMERGENCY, GO TO THE HOSPITAL EMERGENCY ROOM!